GEOMETRIC FORMULAS



Rectangle

Area: $A = \ell w$

Perimeter: $p = 2\ell + 2w$



Square

Area: $A = s^2$

Perimeter: $p = 4s$



Parallelogram

Area: $A = bh$



Triangle

Area: $A = \frac{1}{2}bh$

$m\angle A + m\angle B + m\angle C = 180°$



Trapezoid

Area: $A = \frac{1}{2}h(b_1 + b_2)$



Regular Polygon

Area: $A = \frac{1}{2}ap$



Circle

Area: $A = \pi r^2$

Circumference: $C = \pi d = 2\pi r$



Right Prism

Volume: $V = Bh$

Lateral Area: $LA = ph$

Surface Area: $SA = ph + 2B$



Regular Pyramid

Volume: $V = \frac{1}{3}Bh$

Lateral Area: $LA = \frac{1}{2}ps$

Surface Area: $SA = \frac{1}{2}ps + B$



Right Cylinder

Volume: $V = \pi r^2 h$

Lateral Area: $LA = 2\pi rh$

Surface Area: $SA = 2\pi rh + 2\pi r^2$



Right Cone

Volume: $V = \frac{1}{3}\pi r^2 h$

Lateral Area: $LA = \pi rs$

Surface Area: $SA = \pi rs + \pi r^2$



Sphere

Volume: $V = \frac{4}{3}\pi r^3$

Surface Area: $SA = 4\pi r^2$

$\sec\theta = \frac{r}{x} = \frac{hyp}{adj}$ $\cot\theta = \frac{x}{y} = \frac{adj}{opp}$

$\tan\theta = \frac{y}{x} = \frac{opp}{adj}$ $\csc\theta = \frac{r}{y} = \frac{hyp}{opp}$

$\sin\theta = \frac{y}{r} = \frac{opp}{hyp}$ $\cos\theta = \frac{x}{r} = \frac{adj}{hyp}$

distance $= \sqrt{(x_2-x_1)^2 + (y_2-y_1)^2}$

midpoint $\left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right)$

30-60-90 rt. Δ

shorter leg $= \frac{hyp}{2}$

longer leg $= \sqrt{3}$ (shorter leg)

45-45-90 rt. Δ

hyp $= \sqrt{2}$ (leg)

$ax^2 + bx + c = 0$

$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

FORMULAS FROM GEOMETRY

Triangle

$h = a \sin \theta$

Area $= \frac{1}{2}bh$

(Law of Cosines)

$c^2 = a^2 + b^2 - 2ab \cos \theta$



Sector of Circular Ring

(p = average radius,

w = width of ring,

θ in radians)

Area $= \theta pw$



Right Triangle

(Pythagorean Theorem)

$c^2 = a^2 + b^2$



Ellipse

Area $= \pi ab$

Circumference $\approx 2\pi\sqrt{\frac{a^2 + b^2}{2}}$



Equilateral Triangle

$h = \frac{\sqrt{3}s}{2}$

Area $= \frac{\sqrt{3}s^2}{4}$



Cone

(A = area of base)

Volume $= \frac{Ah}{3}$

Parallelogram

Area $= bh$



Right Circular Cone

Volume $= \frac{\pi r^2 h}{3}$

Lateral Surface Area $= \pi r\sqrt{r^2 + h^2}$



Trapezoid

Area $= \frac{h}{2}(a + b)$



Frustum of Right Circular Cone

Volume $= \frac{\pi(r^2 + rR + R^2)h}{3}$

Lateral Surface Area $= \pi s(R + r)$



Circle

Area $= \pi r^2$

Circumference $= 2\pi r$

Right Circular Cylinder

Volume $= \pi r^2 h$

Lateral Surface Area $= 2\pi rh$



Sector of Circle

(θ in radians)

Area $= \frac{\theta r^2}{2}$

$s = r\theta$



Sphere

Volume $= \frac{4}{3}\pi r^3$

Surface Area $= 4\pi r^2$



Circular Ring

(p = average radius,

w = width of ring)

Area $= \pi(R^2 - r^2)$

$= 2\pi pw$



Wedge

(A = area of upper face,

B = area of base)

$A = B \sec \theta$



Areas Of Quads

NAME	Drawing	FORMULAS	Description
Rectangle		$A=bh$	
Square		$A=s^2$	
Parallelogram		$A=bh$	
Trapezoid		$A=\frac{1}{2}(b_1+b_2)h$	
Kite		$A=\frac{ab}{2}$	
Rhombus		$A=sh$ or $A=\frac{ab}{2}$	
* HERON'S FORMULA *	 $s=\frac{1}{2}(a+b+c)$	$A=\sqrt{s(s-a)(s-b)(s-c)}$	

Trigonometric Functions	Right Triangle Ratios ($0° < \theta < 90°$)
$\sin\theta = \frac{y}{r}$	$\sin\theta = \frac{a}{c} = \frac{\text{length of leg opposite } \theta}{\text{length of hypotenuse}}$
$\cos\theta = \frac{x}{r}$	$\cos\theta = \frac{b}{c} = \frac{\text{length of leg adjacent to } \theta}{\text{length of hypotenuse}}$
$\tan\theta = \frac{y}{x}$	$\tan\theta = \frac{a}{b} = \frac{\text{length of leg opposite } \theta}{\text{length of leg adjacent to } \theta}$
$\csc\theta = \frac{r}{y}$	$\csc\theta = \frac{c}{a} = \frac{\text{length of hypotenuse}}{\text{length of leg opposite } \theta}$
$\sec\theta = \frac{r}{x}$	$\sec\theta = \frac{c}{b} = \frac{\text{length of hypotenuse}}{\text{length of leg adjacent to } \theta}$
$\cot\theta = \frac{x}{y}$	$\cot\theta = \frac{b}{a} = \frac{\text{length of leg adjacent to } \theta}{\text{length of leg opposite } \theta}$



Capital letters are usually used to represent the angles of triangles, or their measures. Lowercase letters refer to the sides opposite their respective angles, or to their measures. The right triangle ratios can be used to *solve* a right triangle, that is, to find the unknown measures of the sides and angles.

EXAMPLE 1 **Solve right triangle ABC if $b = 32$, $\angle A = 25°$, and $\angle C = 90°$. Fin and c to the nearest unit.**

To find a, use $\tan 25°$.

$$\tan 25° = \frac{a}{32} \qquad \tan A = \frac{a}{b}$$

$$a = 32 \tan 25° \qquad \text{Calculation-ready form}$$

$$a = 15 \qquad \text{To the nearest unit}$$



To find c, use $\cos 25°$.

$$\cos 25° = \frac{32}{c} \qquad \cos A = \frac{b}{c}$$

$$c = \frac{32}{\cos 25°} = 35 \qquad \text{To the nearest unit}$$

Since angles A and B are complementary, $\angle B = 90° - 25° = 65°$.

A **significant digit** is any nonzero digit or any zero that serves a purpose other than to locate the decimal point. Consider the following examples:

0.00*304*	Three significant digits
29.40	Four significant digits
[illegible]	[illegible] significant digits

*Keep For Pre-Calculus

TRIGONOMETRY

Unit Circle

Definition of the Six Trigonometric Functions

Right triangle definitions, where $0 < \theta < \pi/2$.



$$\sin\theta = \frac{\text{opp}}{\text{hyp}} \quad \csc\theta = \frac{\text{hyp}}{\text{opp}}$$

$$\cos\theta = \frac{\text{adj}}{\text{hyp}} \quad \sec\theta = \frac{\text{hyp}}{\text{adj}}$$

$$\tan\theta = \frac{\text{opp}}{\text{adj}} \quad \cot\theta = \frac{\text{adj}}{\text{opp}}$$

Circular function definitions, where θ is any angle.



$$\sin\theta = \frac{y}{r} \quad \csc\theta = \frac{r}{y}$$

$$\cos\theta = \frac{x}{r} \quad \sec\theta = \frac{r}{x}$$

$$\tan\theta = \frac{y}{x} \quad \cot\theta = \frac{x}{y}$$



Reciprocal Identities

$$\sin x = \frac{1}{\csc x} \quad \sec x = \frac{1}{\cos x} \quad \tan x = \frac{1}{\cot x}$$

$$\csc x = \frac{1}{\sin x} \quad \cos x = \frac{1}{\sec x} \quad \cot x = \frac{1}{\tan x}$$

Tangent and Cotangent Identities

$$\tan x = \frac{\sin x}{\cos x} \quad \cot x = \frac{\cos x}{\sin x}$$

Pythagorean Identities

$$\sin^2 x + \cos^2 x = 1$$

$$1 + \tan^2 x = \sec^2 x \qquad 1 + \cot^2 x = \csc^2 x$$

Cofunction Identities

$$\sin\left(\frac{\pi}{2} - x\right) = \cos x \quad \cos\left(\frac{\pi}{2} - x\right) = \sin x$$

$$\csc\left(\frac{\pi}{2} - x\right) = \sec x \quad \tan\left(\frac{\pi}{2} - x\right) = \cot x$$

$$\sec\left(\frac{\pi}{2} - x\right) = \csc x \quad \cot\left(\frac{\pi}{2} - x\right) = \tan x$$

Reduction Formulas

$$\sin(-x) = -\sin x \quad \cos(-x) = \cos x$$

$$\csc(-x) = -\csc x \quad \tan(-x) = -\tan x$$

$$\sec(-x) = \sec x \quad \cot(-x) = -\cot x$$

Sum and Difference Formulas

$$\sin(u \pm v) = \sin u \cos v \pm \cos u \sin v$$

$$\cos(u \pm v) = \cos u \cos v \mp \sin u \sin v$$

$$\tan(u \pm v) = \frac{\tan u \pm \tan v}{1 \mp \tan u \tan v}$$

Double-Angle Formulas

$$\sin 2u = 2 \sin u \cos u$$

$$\cos 2u = \cos^2 u - \sin^2 u = 2\cos^2 u - 1 = 1 - 2\sin^2 u$$

$$\tan 2u = \frac{2 \tan u}{1 - \tan^2 u}$$

Power-Reducing Formulas

$$\sin^2 u = \frac{1 - \cos 2u}{2}$$

$$\cos^2 u = \frac{1 + \cos 2u}{2}$$

$$\tan^2 u = \frac{1 - \cos 2u}{1 + \cos 2u}$$

Sum-to-Product Formulas

$$\sin u + \sin v = 2\sin\left(\frac{u+v}{2}\right)\cos\left(\frac{u-v}{2}\right)$$

$$\sin u - \sin v = 2\cos\left(\frac{u+v}{2}\right)\sin\left(\frac{u-v}{2}\right)$$

$$\cos u + \cos v = 2\cos\left(\frac{u+v}{2}\right)\cos\left(\frac{u-v}{2}\right)$$

$$\cos u - \cos v = -2\sin\left(\frac{u+v}{2}\right)\sin\left(\frac{u-v}{2}\right)$$

Product-to-Sum Formulas

$$\sin u \sin v = \frac{1}{2}[\cos(u - v) - \cos(u + v)]$$

$$\cos u \cos v = \frac{1}{2}[\cos(u - v) + \cos(u + v)]$$

$$\sin u \cos v = \frac{1}{2}[\sin(u + v) + \sin(u - v)]$$

$$\cos u \sin v = \frac{1}{2}[\sin(u + v) - \sin(u - v)]$$

FINITE DIFFERENCE RULE

1. QUADRATIC EXPRESSION (QUADRATIC PATTERNS)

$$an^2 + bn + c$$

TERM	VALUE	D1	D2
0	c=		
1	6	a + b=	a=(D2)/2
2	28		a=(D2)/2
3	65		a=(D2)/2
4	117		a=(D2)/2
5	184		a=(D2)/2
6	266		a=(D2)/2
n			

2.LINEAR EXPRESSION (LINEAR PATTERNS)

$$an + b$$

TERM	VALUE	D1	
0	b =		
1	-1	a= D1	
2	-6	a= D1	
3	-11	a= D1	
4	-16	a= D1	
5	-21	a= D1	
6	-26	a= D1	
n			

;QUADRATIC PARENT FUNCTION :

NAME: _____ ; **PERIOD:** ____ .



x	y
1	1
2	4
3	9
-1	1
-2	4
-3	9

Quadratic Functions

1) $y = ax^2 + bx + c$
2) Non-linear funtions
3) $y = x^2$ is Parent Function
4)Maximum value at vertex { ∩ } or Minimum value at vertex { U }
5) Line of symmetry (axis of symmetry)
6) Root(s) or x-intercept (s) or Solution(s)

;QUADRATIC FUNCTION GRAPH COMPONENTS:

NAME: $y = ax^2 + bx + c$;

If a < 0, then you have a Maximum Value graph. If a >0, Then you have Minimum Value graph. If a = 0, then you have a linear graph not a quadratic anymore. c is always the y-intercept.



;QUADRATIC GRAPH COMPONENTS EQUATIONS:

NAME: $y = ax^2 + bx + c$;

If $a < 0$, then you have a Maximum Value graph. If $a > 0$, Then you have Minimum Value graph. If $a = 0$, then you have a linear graph not a quadratic anymore. c is always the y-intercept.



;QUADRATIC FUNCTION GRAPH PROPERTIES:

NAME: $y = ax^2 + bx + c$;

If $a < 0$, then you have a Maximum Value graph. If $a > 0$, Then you have Minimum Value graph. If $a = 0$, then you have a linear graph not a quadratic anymore. c is always the y-intercept.

0 Solutions = imaginary roots



ALGEBRA

Factors and Zeros of Polynomials

Let $p(x) = a_n x^n + a_{n-1}x^{n-1} + \cdots + a_1 x + a_0$ be a polynomial. If $p(a) = 0$, then a is a *zero* of the polynomial and a solution of the equation $p(x) = 0$. Furthermore, $(x - a)$ is a *factor* of the polynomial.

Fundamental Theorem of Algebra

An nth degree polynomial has n (not necessarily distinct) zeros. Although all of these zeros may be imaginary, a real polynomial of odd degree must have at least one real zero.

Quadratic Formula

If $p(x) = ax^2 + bx + c$, and $0 \le b^2 - 4ac$, then the real zeros of p are $x = \left(-b \pm \sqrt{b^2 - 4ac}\right)/2a$.

Special Factors

$x^2 - a^2 = (x - a)(x + a)$

$x^3 - a^3 = (x - a)(x^2 + ax + a^2)$

$x^3 + a^3 = (x + a)(x^2 - ax + a^2)$

$x^4 - a^4 = (x^2 - a^2)(x^2 + a^2)$

Binomial Theorem

$(x + y)^2 = x^2 + 2xy + y^2$

$(x - y)^2 = x^2 - 2xy + y^2$

$(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$

$(x - y)^3 = x^3 - 3x^2y + 3xy^2 - y^3$

$(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

$(x - y)^4 = x^4 - 4x^3y + 6x^2y^2 - 4xy^3 + y^4$

$$(x + y)^n = x^n + nx^{n-1}y + \frac{n(n-1)}{2!}x^{n-2}y^2 + \cdots + nxy^{n-1} + y^n$$

$$(x - y)^n = x^n - nx^{n-1}y + \frac{n(n-1)}{2!}x^{n-2}y^2 - \cdots \pm nxy^{n-1} \mp y^n$$

Rational Zero Theorem

If $p(x) = a_n x^n + a_{n-1}x^{n-1} + \cdots + a_1 x + a_0$ has integer coefficients, then every *rational zero* of p is of the form $x = r/s$, where r is a factor of a_0 and s is a factor of a_n.

Factoring by Grouping

$acx^3 + adx^2 + bcx + bd = ax^2(cx + d) + b(cx + d) = (ax^2 + b)(cx + d)$

Arithmetic Operations

$ab + ac = a(b + c)$

$\dfrac{a}{b} + \dfrac{c}{d} = \dfrac{ad + bc}{bd}$

$\dfrac{a + b}{c} = \dfrac{a}{c} + \dfrac{b}{c}$

$\dfrac{\left(\frac{a}{b}\right)}{\left(\frac{c}{d}\right)} = \left(\dfrac{a}{b}\right)\left(\dfrac{d}{c}\right) = \dfrac{ad}{bc}$

$\dfrac{\left(\frac{a}{b}\right)}{c} = \dfrac{a}{bc}$

$\dfrac{a}{\left(\frac{b}{c}\right)} = \dfrac{ac}{b}$

$a\left(\dfrac{b}{c}\right) = \dfrac{ab}{c}$

$\dfrac{a - b}{c - d} = \dfrac{b - a}{d - c}$

$\dfrac{ab + ac}{a} = b + c$

Exponents and Radicals

$a^0 = 1, \quad a \neq 0$

$(ab)^x = a^x b^x$

$a^x a^y = a^{x+y}$

$\sqrt{a} = a^{1/2}$

$\dfrac{a^x}{a^y} = a^{x-y}$

$\sqrt[n]{a} = a^{1/n}$

$\left(\dfrac{a}{b}\right)^x = \dfrac{a^x}{b^x}$

$\sqrt[n]{a^m} = a^{m/n}$

$a^{-x} = \dfrac{1}{a^x}$

$\sqrt[n]{ab} = \sqrt[n]{a}\,\sqrt[n]{b}$

$(a^x)^y = a^{xy}$

$\sqrt[n]{\dfrac{a}{b}} = \dfrac{\sqrt[n]{a}}{\sqrt[n]{b}}$

Notes: Exponentiation

Definitions:

An **exponential function** is a function whose general equation is $y = a \cdot b^x$ where *a* and *b* stand for constants, b is positive and *x* and *y* are independent and dependent variables.

Exponentiation for positive integer exponent: (when *n* is positive) x^n means the product of *n x*'s.

In the expression x^n,

x is called the *base,*
n is called the *exponent*
x^n is called a *power.*

Example 1: $-x^3 = -(x)(x)(x)$
$(-x)^3 = (-x)(-x)(-x)$

Example 2: $4x^4 = 4(x)(x)(x)(x)$

$(4x)^4 = (4x)(4x)(4x)(4x)$

Rule: If the *base* contains more than one symbol, then the *base* must be placed in <u>parentheses</u>.

Properties of Exponentiation:

1. *Product of two powers with equal bases:*

$$x^a \cdot x^b = x^{a+b}$$

2. *Quotient of two powers with equal bases:*

$$\frac{x^a}{x^b} = x^{a-b}$$

3. *Power of a power:*

$$(x^a)^b = x^{a*b}$$

Log Rules

$$a = b^{c} \Rightarrow \log_b a = c$$

Standard Form — Base — Exponent

standard — base — Exponent

① $a = b^{c} \Rightarrow \log_b a = c$

② $\log_b MN \Rightarrow \log_b M + \log_b N$

③ $\log_b \frac{M}{N} \Rightarrow \log_b M - \log_b N$

④ $\log_b N^{a} \Rightarrow a \log_b N$

⑤ $\log_b a = \frac{\log a}{\log b}$

eg: $\log_5 3 = \frac{\log 3}{\log 5}$ use calculator Now.

CONICS

PARABOLA

$$y-k=a(x-h)^2$$

$$x-h=a(y-k)^2$$

$$a=\frac{1}{4p}$$





CIRCLE

$$(x-h)^2+(y-k)^2=r^2$$



ELLIPSE

$$\frac{(x-h)^2}{a^2}+\frac{(y-k)^2}{b^2}=1$$





HYPERBOLA

$$\frac{(x-h)^2}{a^2}-\frac{(y-k)^2}{b^2}=1 \text{ or } \frac{(y-k)^2}{b^2}-\frac{(x-h)^2}{a^2}=1$$





SPECIAL HYPERBOLAS

$$xy=k \text{ or } xy=-k$$





GRAPHS OF COMMON FUNCTIONS

PARENT FUNCTIONS



Constant Function



Identity Function



Absolute Value Function



Square Root Function



Squaring Function



Cubing Function



Exponential Function



Logarithmic Function

Rational Function

$f(x) = \frac{1}{x}$



SYMMETRY



y-Axis Symmetry



x-Axis Symmetry



Origin Symmetry

Transformations for $y = f(x)$

$y = f(x) + k$	vertical shift k units up
$y = f(x) - k$	vertical shift k units down
$y = f(x - h)$	horizontal shift h units right
$y = f(x + h)$	horizontal shift h units left
$y = -f(x)$	reflected in the x-axis
$y = f(-x)$	reflected in the y-axis
$x = f(y)$	reflected in the line $y = x$
$y = \lvert f(x) \rvert$	unchanged when $f(x) > 0$; reflected in the x-axis when $f(x) < 0$
$y = f(\lvert x \rvert)$	unchanged when $x > 0$; reflected in the y-axis when $x < 0$

multiply "y" values by "a"

$y = af(x), \lvert a \rvert > 1$	stretched vertically
$y = af(x), 0 < \lvert a \rvert < 1$	shrunk vertically

multiply "x" values by $\frac{1}{b}$

$y = f(bx), \lvert b \rvert > 1$	shrunk horizontally
$y = f(bx), 0 < \lvert b \rvert < 1$	stretched horizontally

Vertical stretch or shrink; reflection across x-axis

Vertical shift

$$y = a\,f(b(x+c))+d$$

Horizontal stretch or shrink; reflection across y-axis

Horizontal shift

Interval notation is used in Algebra II for many topics.

1. Symbols:
   a. ∞ infinity
   b. $-\infty$ negative infinity
   c. [or] the value is included
   d. (or) the value is not included
   e. $\cup$ union or joining together two sets of numbers
2. Infinity <u>always</u> using soft brackets.

Interval Notation	*Set Builder Notation*	*Inequality Notation*	*Graph of Inequality*
$(-\infty, a]$	$\{x \mid x \le a\}$	$x \le a$	
$(-\infty, a)$	$\{x \mid x < a\}$	$x < a$	
$[b, \infty)$	$\{x \mid x \ge b\}$	$x \ge b$	
(b, ∞)	$\{x \mid x > b\}$	$x > b$	
$[a, b]$	$\{x \mid a \le x \le b\}$	$a \le x \le b$	
$[a, b)$	$\{x \mid a \le x < b\}$	$a \le x < b$	
$(a, b]$	$\{x \mid a < x \le b\}$	$a < x \le b$	
(a, b)	$\{x \mid a < x < b\}$	$a < x < b$	
$(-\infty, a] \cup (b, \infty)$	$\{x \mid x \le a \text{ or } x > b\}$	$x \le a, x > b$